

Constitutive Theory

Common constitutive laws:

Newtonian fluid: $\underline{\underline{\sigma}} = -p \underline{\underline{I}} + \eta (\nabla \underline{v} + \nabla \underline{v}^T)$

$$p = -\frac{1}{3} \text{tr}(\underline{\underline{\sigma}}) \quad \eta = \text{viscosity} \quad \underline{v} = \text{velocity}$$

Linear elastic solid: $\underline{\underline{\sigma}} = \lambda \nabla \cdot \underline{u} \underline{\underline{I}} + \mu (\nabla \underline{u} + \nabla \underline{u}^T)$

$$\lambda, \mu = \text{Lame parameters} \quad \underline{u} = \text{displacement}$$

Both derive from the functional form

$$\underline{\underline{G}}(\underline{\underline{E}}) = \lambda \text{tr}(\underline{\underline{E}}) \underline{\underline{I}} + 2\mu \text{sym}(\underline{\underline{E}})$$

Newtonian fluid: $\underline{\underline{E}} = \nabla \underline{v}$

Linear elastic solid: $\underline{\underline{E}} = \nabla \underline{u}$

remember $\nabla \cdot \underline{a} = \text{tr}(\nabla \underline{a})$

⇒ direct for lin. elastic solid

for fluid there is a complication due to incompressibility!

Why do const. relations have this form?

Change of observer

In Lecture 6 we discussed Change in basis

$$\underline{v} = \underline{Q} \underline{v}' \quad \text{and} \quad \underline{S} = \underline{Q} \underline{S}' \underline{Q}^T$$


where \underline{Q} is change in basis tensor.

Q is a rotation: 1) orthonormal $\underline{Q} \underline{Q}^T = \underline{Q}^T \underline{Q} = \underline{I}$

$$2) \det(\underline{Q}) = 1$$

Change in basis is passive change of frame.

Active change in frame \rightarrow change in observer



$$\underline{x} = \varphi(\underline{x}, t)$$

$$\underline{x}' = \varphi'(\underline{x}, t)$$

Note: Material ref.

frame is common

Note: We assume both observers

use same clock.

Since change in observer cannot induce a deformation. Two ref. frames must be related by a rigid body motion.

$$\underline{x}' = Q(t) \varphi(\underline{x}, t) + \underline{c}(t) \quad \text{Eulerian transformation}$$

$$\underline{Q} = \text{rotation} \quad \underline{c} = \text{translation}$$

Our description of forces and deformations cannot depend on the observer (objective).

Effect on kinematic quantities

$$\underline{x} = \varphi(\underline{x}, t) \quad \nabla \varphi = \underline{F}$$

$$\underline{x}' = \varphi'(\underline{x}, t) = Q \varphi(\underline{x}, t) + c \quad \nabla \varphi' = \underline{Q} \underline{F} = \underline{F}'$$

Right Cauchy-Green Strain tensor

$$\underline{C}' = \underline{F}'^T \underline{F}' = (\underline{Q} \underline{F})^T (\underline{Q} \underline{F}) = \underline{F}^T \underline{Q}^T \underline{Q} \underline{F} = \underline{F}^T \underline{F} = \underline{C}$$

⇒ not affected by rigid body motion because

it is a material tensor C_{IJ} (naturally objective)

What about spatial tensors?

Axiom of frame indifference

Fields ϕ , $\underline{\omega}$ and $\underline{\underline{S}}$ are called frame indifferent or objective if for all superposed rigid body motions $\underline{x}' = \underline{Q}\underline{x} + \underline{c}$ we have for all spatial fields

$\phi'(\underline{x}', t) = \phi(\underline{x}, t)$	scalar field
$\underline{\omega}'(\underline{x}', t) = \underline{Q} \underline{\omega}(\underline{x}, t)$	vector field
$\underline{\underline{S}}'(\underline{x}', t) = \underline{Q} \underline{\underline{S}}(\underline{x}, t) \underline{Q}^T$	tensor field

\Rightarrow from Lecture 6.

Is spatial velocity gradient objective?

From Lecture 16: $\underline{\underline{L}} = \nabla_{\underline{x}} \underline{v} = \underline{\underline{\dot{F}}} \underline{\underline{F}}^{-1}$

$$\underline{\underline{F}}' = \underline{Q} \underline{\underline{F}} \quad \underline{\underline{L}}' = \nabla_{\underline{x}'} \underline{v}' = \underline{\underline{\dot{F}}}' \underline{\underline{F}}'^{-1}$$

$$\underline{\underline{\dot{F}}}' = \frac{d}{dt} (\underline{Q} \underline{\underline{F}}) = \underline{Q} \underline{\underline{\dot{F}}} + \underline{\underline{\dot{Q}}} \underline{\underline{F}}$$

$$\underline{\underline{F}}'^{-1} = (\underline{Q} \underline{\underline{F}})^{-1} = \underline{\underline{F}}^{-1} \underline{Q}^{-1} = \underline{\underline{F}}^{-1} \underline{Q}^T$$

$$\underline{\underline{L}}' = \underline{\underline{\dot{F}}}' \underline{\underline{F}}'^{-1} = (\underline{Q} \underline{\underline{\dot{F}}} + \underline{\underline{\dot{Q}}} \underline{\underline{F}}) \underline{\underline{F}}^{-1} \underline{Q}^T$$

$$= \underline{Q} \underline{\underline{\dot{F}}} \underline{\underline{F}}^{-1} \underline{Q}^T + \underline{\underline{\dot{Q}}} \underline{\underline{F}} \underline{\underline{F}}^{-1} \underline{Q}^T = \underline{Q} \underline{\underline{L}} \underline{Q}^T + \underline{\underline{\dot{Q}}} \underline{Q}^T$$

$$\Rightarrow \underline{\underline{\ell}}' = \underline{\underline{Q}} \underline{\underline{\ell}} \underline{\underline{Q}}^T + \underline{\underline{\dot{Q}}} \underline{\underline{Q}}^T \quad \underline{\underline{\text{not objective!}}}$$

that is why $\nabla_x \underline{v}$ is not used in constitutive laws

The "non-objective" term is $\underline{\underline{\Omega}} = \underline{\underline{\dot{Q}}} \underline{\underline{Q}}^T$

it represents rigid body angular velocity between observers. *see HW9*

Show $\underline{\underline{\Omega}} = -\underline{\underline{\Omega}}^T$ skew-symmetric

Non-objective part of $\underline{\underline{\ell}} = \nabla_x \underline{v}$ is skew-sym.

\Rightarrow simply take symmetric part of $\underline{\underline{\ell}}$!

$$\underline{\underline{d}} = \text{sym}(\underline{\underline{\ell}}) = \frac{1}{2} (\nabla_x \underline{v} + \nabla_x \underline{v}^T)$$

rate of strain tensor is objective

\Rightarrow used in constitutive laws

Note that velocity itself

Material frame indifferent functions

Fields: $\phi(\underline{x}, t)$ scalar

$\underline{w}(\underline{x}, t)$ vector

$\underline{\underline{s}}(\underline{x}, t)$ tensor

fields because they depend on \underline{x} .

Constitutive functions are not fields but they depend on fields as input.

internal energy: $u(\underline{x}, t) = \hat{u}(\rho(\underline{x}, t), \theta(\underline{x}, t))$
output field \uparrow constitutive function \uparrow input fields

heat flow: $\underline{q}(\underline{x}, t) = \hat{\underline{q}}(\theta(\underline{x}, t))$

Cauchy stress: $\underline{\underline{s}}(\underline{x}, t) = \hat{\underline{\underline{s}}}(\rho(\underline{x}, t), \theta(\underline{x}, t), \underline{d}(\underline{x}, t))$

Constitutive functions: $\hat{u}(\rho, \theta)$, $\hat{\underline{q}}(\theta)$, $\hat{\underline{\underline{s}}}(\rho, \theta, \underline{d})$

As such constitutive functions are not directly dependent on frame but their input fields are.

Consider frames $\{\underline{e}_i\}$ and $\{\underline{e}'_i\}$ then to be frame indifference requires

$$\hat{\underline{\underline{\sigma}}}(\rho', \theta', \underline{d}') = \underline{Q} \hat{\underline{\underline{\sigma}}}(\rho, \theta, \underline{d}) \underline{Q}^T$$

substituting $\underline{d}' = \underline{Q} \underline{d} \underline{Q}^T$

$$\hat{\underline{\underline{\sigma}}}(\rho', \theta', \underline{Q} \underline{d} \underline{Q}^T) = \underline{Q} \hat{\underline{\underline{\sigma}}}(\rho, \theta, \underline{d}) \underline{Q}^T$$

⇒ both input & output of constitutive function $\hat{\underline{\underline{\sigma}}}$ must be frame invariant

Isotropic functions

Functions that are frame invariant are called isotropic. Consider the following

$\hat{\phi}$ = scalar fun. $\hat{\underline{w}}$ = vector fun. $\hat{\underline{\underline{s}}}$ = tensor fun.

θ = scalar \underline{v} = vector $\underline{\underline{s}}$ = tensor

Then for two frames related by rigid body rotation $\underline{\underline{Q}}$ we have following isotropic functions:

$$\hat{\phi}(\theta) = \hat{\phi}(\theta) \quad \hat{\phi}(\underline{\underline{Q}}\underline{v}) = \hat{\phi}(\underline{v}) \quad \hat{\phi}(\underline{\underline{Q}}\underline{\underline{s}}\underline{\underline{Q}}^T) = \hat{\phi}(\underline{\underline{s}})$$

$$\hat{\underline{w}}(\theta) = \underline{\underline{Q}}\hat{\underline{w}}(\theta) \quad \hat{\underline{w}}(\underline{\underline{Q}}\underline{v}) = \underline{\underline{Q}}\hat{\underline{w}}(\underline{v}) \quad \hat{\underline{w}}(\underline{\underline{Q}}\underline{\underline{s}}\underline{\underline{Q}}^T) = \underline{\underline{Q}}\hat{\underline{w}}(\underline{\underline{s}})$$

$$\hat{\underline{\underline{s}}}(\theta) = \underline{\underline{Q}}\hat{\underline{\underline{s}}}(\theta)\underline{\underline{Q}}^T \quad \hat{\underline{\underline{s}}}(\underline{\underline{Q}}\underline{v}) = \underline{\underline{Q}}\hat{\underline{\underline{s}}}(\underline{v})\underline{\underline{Q}}^T \quad \hat{\underline{\underline{s}}}(\underline{\underline{Q}}\underline{\underline{s}}\underline{\underline{Q}}^T) = \underline{\underline{Q}}\hat{\underline{\underline{s}}}(\underline{\underline{s}})\underline{\underline{Q}}^T$$

Examples:

1) $\hat{\phi}(\underline{\underline{s}}) = \det(\underline{\underline{s}})$

$$\hat{\phi}(\underline{\underline{Q}}\underline{\underline{s}}\underline{\underline{Q}}^T) = \det(\underline{\underline{Q}}\underline{\underline{s}}\underline{\underline{Q}}^T) = \det(\underline{\underline{Q}})\det(\underline{\underline{s}})\det(\underline{\underline{Q}}^T) = \det(\underline{\underline{s}}) \checkmark$$

2) $\hat{\underline{w}}(\underline{v}, \underline{\underline{A}}) = \underline{\underline{A}}\underline{v}$

$$\hat{\underline{w}}(\underline{\underline{Q}}\underline{v}, \underline{\underline{Q}}\underline{\underline{A}}\underline{\underline{Q}}^T) = \underline{\underline{Q}}\underline{\underline{A}}\underline{\underline{Q}}^T \underline{\underline{Q}}\underline{v} = \underline{\underline{Q}}\underline{\underline{A}}\underline{v} = \underline{\underline{Q}}\hat{\underline{w}}(\underline{v}, \underline{\underline{A}}) \checkmark$$

Isotropic material: stress/strain principal directions

Objectivity \Rightarrow isotropic function

fluids: $\underline{\underline{Q}} \hat{\underline{\underline{\sigma}}}(\underline{\underline{d}}) \underline{\underline{Q}}^T = \hat{\underline{\underline{\sigma}}}(\underline{\underline{Q}} \underline{\underline{d}} \underline{\underline{Q}}^T)$ rate of strain

solids: $\underline{\underline{Q}} \hat{\underline{\underline{\sigma}}}(\underline{\underline{\epsilon}}) \underline{\underline{Q}}^T = \hat{\underline{\underline{\sigma}}}(\underline{\underline{Q}} \underline{\underline{\epsilon}} \underline{\underline{Q}}^T)$ strain

generic: $\underline{\underline{Q}} \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{Q}}^T = \underline{\underline{G}}(\underline{\underline{Q}} \underline{\underline{A}} \underline{\underline{Q}}^T)$

Since $\underline{\underline{\sigma}} = \underline{\underline{\sigma}}^T$ and $\underline{\underline{d}} = \underline{\underline{d}}^T$ ($\underline{\underline{\epsilon}} = \underline{\underline{\epsilon}}^T$) they can all be written in spectral decomposition.

$$\underline{\underline{S}} = \underline{\underline{S}}^T \Rightarrow \underline{\underline{S}} = \sum_{i=1}^3 \alpha_i \underline{\underline{v}}_i \otimes \underline{\underline{v}}_i$$

where $\underline{\underline{S}} \underline{\underline{v}}_i = \alpha_i \underline{\underline{v}}_i$

Q: How are $\underline{\underline{v}}_i$'s of $\underline{\underline{\sigma}}$ and $\underline{\underline{d}}/\underline{\underline{\epsilon}}$ related?


\Rightarrow same eigenvectors!

$$\underline{\underline{A}} \underline{\underline{v}}_i = \alpha_i \underline{\underline{v}}_i \Rightarrow \underline{\underline{G}}(\underline{\underline{A}}) \omega_i = \omega_i \underline{\underline{v}}_i$$

Can be shown with reflections & projections

Projection: $\underline{\underline{P}}_n = \underline{\underline{n}} \otimes \underline{\underline{n}}$

Reflection: $\underline{\underline{R}}_n = \underline{\underline{I}} - 2 \underline{\underline{n}} \otimes \underline{\underline{n}}$



Note: $\underline{\underline{R}}_n \underline{n} = -\underline{n}$

$\underline{\underline{R}}_n \underline{a} = \underline{a}$ if $\underline{a} \cdot \underline{n} = 0$ $\underline{a} \perp \underline{n}$

\Rightarrow reflections help to detect colinear vectors.

If $\underline{n} = \underline{v}_1$ one eigenvectors

$\Rightarrow \underline{\underline{R}}_{\underline{v}_1} \underline{v}_1 = -\underline{v}_1$ but $\underline{\underline{R}}_{\underline{v}_1} \underline{v}_2 = \underline{v}_2$ & $\underline{\underline{R}}_{\underline{v}_1} \underline{v}_3 = \underline{v}_3$

Step 1: $\underline{\underline{R}}_{\underline{v}_1} \underline{A} \underline{\underline{R}}_{\underline{v}_1}^T = \underline{A}$ $\underline{A} = \underline{A}^T$

$= \underline{\underline{R}}_{\underline{v}_1} \left(\sum_{i=1}^3 \alpha_i (\underline{v}_i \otimes \underline{v}_i) \right) \underline{\underline{R}}_{\underline{v}_1}^T$

$= \sum_{i=1}^3 \alpha_i \underline{\underline{R}}_{\underline{v}_1} (\underline{v}_i \otimes \underline{v}_i) \underline{\underline{R}}_{\underline{v}_1}^T$

use identities: $\underline{\underline{S}} (\underline{a} \otimes \underline{b}) = (\underline{\underline{S}} \underline{a}) \otimes \underline{b}$

$(\underline{a} \otimes \underline{b}) \underline{\underline{S}} = \underline{a} \otimes (\underline{\underline{S}}^T \underline{b})$

$\Rightarrow \underline{\underline{R}}_{\underline{v}_1} \underline{A} \underline{\underline{R}}_{\underline{v}_1}^T = \sum_{i=1}^3 \alpha_i (\underline{\underline{R}}_{\underline{v}_1} \underline{v}_i) \otimes (\underline{\underline{R}}_{\underline{v}_1} \underline{v}_i)$

$= \alpha_1 (-\underline{v}_1) \otimes (-\underline{v}_1) + \alpha_2 \underline{v}_2 \otimes \underline{v}_2 + \alpha_3 \underline{v}_3 \otimes \underline{v}_3$

$= \sum_{i=1}^3 \alpha_i \underline{v}_i \otimes \underline{v}_i = \underline{A}$

$\Rightarrow \underline{\underline{R}}_{\underline{v}_1} \underline{d} \underline{\underline{R}}_{\underline{v}_1}^T = \underline{d}$

$\underline{\underline{R}}_{\underline{v}_1} \underline{\underline{e}} \underline{\underline{R}}_{\underline{v}_1}^T = \underline{\underline{e}}$

Step 2: $\underline{\underline{R}}_{v_i} \underline{\underline{G}}(\underline{\underline{A}}) = \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{R}}_{v_i}$ commute

isotropic material: $\underline{\underline{Q}} \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{Q}}^T = \underline{\underline{G}}(\underline{\underline{Q}} \underline{\underline{A}} \underline{\underline{Q}}^T)$

$\underline{\underline{Q}}$ = orthogonal (rotation or reflection)

$\underline{\underline{Q}} = \underline{\underline{R}}_{v_i}$: $\underline{\underline{R}}_{v_i} \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{R}}_{v_i}^T = \underline{\underline{G}}(\underline{\underline{R}}_{v_i} \underline{\underline{A}} \underline{\underline{R}}_{v_i}^T)$

$$= \underline{\underline{G}}(\underline{\underline{A}})$$

$$\underline{\underline{R}}_{v_i} \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{R}}_{v_i}^T \underline{\underline{R}}_{v_i} \underline{\underline{R}}_{v_i}^T = \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{R}}_{v_i}$$

$$\Rightarrow \underline{\underline{R}}_{v_i} \underline{\underline{G}}(\underline{\underline{A}}) = \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{R}}_{v_i}$$

Step 3: $\underline{\underline{A}} \underline{\underline{v}}_i = \alpha_i \underline{\underline{v}}_i \Rightarrow \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{v}}_i = \omega_i \underline{\underline{v}}_i$

$$\underline{\underline{R}}_{v_i} \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{v}}_i = \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{R}}_{v_i} \underline{\underline{v}}_i$$

$$\underline{\underline{R}}_{v_i} \underbrace{\underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{v}}_i}_{\underline{\underline{a}}} = - \underbrace{\underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{v}}_i}_{\underline{\underline{a}}} \Rightarrow \underline{\underline{R}}_{v_i} \underline{\underline{a}} = -\underline{\underline{a}}$$

since $\underline{\underline{R}}_{v_i}$ is reflection $\Rightarrow \underline{\underline{a}} = \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{v}}_i \parallel \underline{\underline{v}}_i$

$\underline{\underline{v}}_i$ is only stretched by $\underline{\underline{G}}(\underline{\underline{A}}) \Rightarrow$ an eigenvector of $\underline{\underline{G}}$

$$\Rightarrow \underline{\underline{G}}(\underline{\underline{A}}) \underline{\underline{v}}_i = \omega_i \underline{\underline{v}}_i$$

principal directions of stress and strain are same

(for isotropic material)

Proof of Representation Thm (linear, isotropic)

Note: $\underline{\underline{A}} = \sum_{i=1}^3 \alpha_i \underbrace{\underline{v}_i \otimes \underline{v}_i}_{\underline{\underline{P}}_{v_i}} = \sum_{i=1}^3 \alpha_i \underline{\underline{P}}_{v_i}$

where $\underline{\underline{P}}_{v_i} = \underline{v}_i \otimes \underline{v}_i$ projection tensors of

Spectral decomposition of stress response

$$\underline{\underline{G}}(\underline{\underline{A}}) = \underline{\underline{G}}\left(\sum_{i=1}^3 \alpha_i \underline{\underline{P}}_{v_i}\right) = \sum_{i=1}^3 \alpha_i \underline{\underline{G}}(\underline{\underline{P}}_{v_i})$$

⇒ total stress response is sum of stress


response in principal directions $\underline{\underline{G}}(\underline{\underline{P}}_{v_i})$

Eigenproblem for Projection tensors

$$\underline{\underline{P}}_n \beta_i = \beta_i v_i \quad (\beta_i, v_i)$$

$$\beta_1 = 1 \quad \beta_2 = \beta_3 = 0$$

$$\Rightarrow \underline{v}_1 = \underline{n}$$



\underline{v}_2 & \underline{v}_3 are any two perpendicular vectors in plane

For any $\underline{\underline{S}} = \underline{\underline{S}}^T$ with $\underline{v}_1, \underline{v}_2, \underline{v}_3$ indep. β_1 & $\beta = \beta_2 = \beta_3$

$$\underline{\underline{S}} = \beta \underline{\underline{I}} + (\beta_1 - \beta) \underline{\underline{P}}_{\underline{v}_1}$$

From spectral decomposition:

$$\begin{aligned}\underline{\underline{E}} &= \beta_1 \underline{\underline{P}}_{\underline{v}_1} + \beta \underline{\underline{P}}_{\underline{v}_2} + \beta \underline{\underline{P}}_{\underline{v}_3} \\ &= \beta_1 \underline{\underline{P}}_{\underline{v}_1} - \beta \underline{\underline{P}}_{\underline{v}_1} + \beta \underline{\underline{P}}_{\underline{v}_1} + \beta \underline{\underline{P}}_{\underline{v}_2} + \beta \underline{\underline{P}}_{\underline{v}_3} \\ &= (\beta_1 - \beta) \underline{\underline{P}}_{\underline{v}_1} + \beta (\underbrace{\underline{\underline{P}}_{\underline{v}_1} + \underline{\underline{P}}_{\underline{v}_2} + \underline{\underline{P}}_{\underline{v}_3}}_{\underline{\underline{I}}}) \\ \underline{\underline{E}} &= \beta \underline{\underline{I}} + (\beta_1 - \beta) \underline{\underline{P}}_{\underline{v}_1}\end{aligned}$$

Apply to $\underline{\underline{G}}(\underline{\underline{P}}_{\underline{v}_i})$ which has same \underline{v}_i 's as $\underline{\underline{P}}_{\underline{v}_i}$

$$\begin{aligned}\underline{\underline{G}}(\underline{\underline{P}}_{\underline{v}_i}) &= \beta \underline{\underline{I}} + (\beta_1 - \beta) \underline{\underline{P}}_{\underline{v}_i} \\ &= \lambda(\underline{v}_i) \underline{\underline{I}} + 2\mu(\underline{v}_i) \underline{\underline{P}}_{\underline{v}_i}\end{aligned}$$

we can show $\lambda(\underline{v}_1) = \lambda(\underline{v}_2) = \lambda(\underline{v}_3) = \lambda = \text{const.}$

see HW $\mu(\underline{v}_1) = \mu(\underline{v}_2) = \mu(\underline{v}_3) = \mu = \text{const.}$

$$\Rightarrow \underline{\underline{G}}(\underline{\underline{P}}_{\underline{v}_i}) = \lambda \underline{\underline{I}} + 2\mu \underline{\underline{P}}_{\underline{v}_i} \quad \text{for any } \underline{v}_i$$

$$\begin{aligned}
 \underline{\underline{G}}(\underline{\underline{A}}) &= \underline{\underline{\left(\sum_{i=1}^3 \alpha_i \underline{\underline{P}}_{v_i} \right)}} = \sum_{i=1}^3 \alpha_i \underline{\underline{P}}_{v_i} \\
 &= \sum_{i=1}^3 \alpha_i (\lambda \underline{\underline{I}} + 2\mu \underline{\underline{P}}_{v_i}) \\
 &= \lambda (\alpha_1 + \alpha_2 + \alpha_3) \underline{\underline{I}} + 2\mu (\alpha_1 \underline{\underline{P}}_{v_1} + \alpha_2 \underline{\underline{P}}_{v_2} + \alpha_3 \underline{\underline{P}}_{v_3}) \\
 &\quad \underbrace{\hspace{10em}}_{\text{tr}(\underline{\underline{A}})} \quad \underbrace{\hspace{10em}}_{\underline{\underline{A}}}
 \end{aligned}$$

Representation for linear isotropic Tensor function

An linear isotropic function $\underline{\underline{G}}(\underline{\underline{A}})$ that maps symmetric tensors $\underline{\underline{E}}$ into symmetric tensors $\underline{\underline{G}}(\underline{\underline{A}})$ must have following form

$$\underline{\underline{G}}(\underline{\underline{A}}) = \lambda \text{tr}(\underline{\underline{A}}) \underline{\underline{I}} + 2\mu \underline{\underline{A}} \quad \underline{\underline{A}} = \underline{\underline{A}}^T$$

where $\lambda, \mu \in \mathbb{R}$ are scalars

In terms of a non-symmetric tensor $\underline{\underline{E}} \neq \underline{\underline{E}}^T$

$$\underline{\underline{G}}(\underline{\underline{E}}) = \lambda \text{tr}(\underline{\underline{E}}) \underline{\underline{I}} + \mu \text{sym}(\underline{\underline{E}})$$

\Rightarrow standard constitutive laws

Linear elasticity: $\underline{\underline{E}} = \nabla \underline{\underline{u}}$

Newtonian fluid: $\underline{\underline{E}} = \nabla \underline{\underline{v}}$

Show λ & μ are independent of \underline{v}_i :

$$|\underline{e}| = |\underline{f}| = 1 \quad \underline{R} \underline{e} = \underline{f} \quad \underline{R} \underline{R}^T = \underline{I} \quad \det(\underline{R}) = -1$$

$$\begin{aligned} \Rightarrow \underline{P}_f &= \underline{f} \otimes \underline{f} = (\underline{R} \underline{e}) \otimes (\underline{R} \underline{e}) = \underline{R} (\underline{e} \otimes \underline{e}) \underline{R}^T = \\ &= \underline{R} \underline{P}_e \underline{R}^T \end{aligned}$$

isotropic: $\hat{\underline{\sigma}}(\underline{R} \underline{P}_e \underline{R}^T) = \underline{R} \hat{\underline{\sigma}}(\underline{P}_e) \underline{R}^T$

$$\hat{\underline{\sigma}}(\underline{P}_f) = \underline{R} \hat{\underline{\sigma}}(\underline{P}_e) \underline{R}^T$$

$$\Rightarrow \underline{R} \hat{\underline{\sigma}}(\underline{P}_e) \underline{R}^T - \hat{\underline{\sigma}}(\underline{P}_f) = \underline{0}$$

substituting: $\hat{\underline{\sigma}}(\underline{P}_f) = \lambda(\underline{f}) \underline{I} + 2\mu(\underline{f}) \underline{P}_f$

$$\hat{\underline{\sigma}}(\underline{P}_e) = \lambda(\underline{e}) \underline{I} + 2\mu(\underline{e}) \underline{P}_e$$

$$[\lambda(\underline{e}) - \lambda(\underline{f})] \underline{I} + 2[\mu(\underline{e}) - \mu(\underline{f})] \underline{P}_f = \underline{0}$$

since \underline{I} and \underline{P}_f are linearly independent

$$\Rightarrow \lambda(\underline{e}) = \lambda(\underline{f}) = \lambda \quad \mu(\underline{e}) = \mu(\underline{f}) = \mu$$

λ & μ are constants

$$\hat{\underline{\sigma}}(\underline{P}_{v_i}) = \lambda \underline{I} + 2\mu \underline{P}_{v_i}$$